

Feed supplement containing live and inactive yeast



- stabilizes the rumen, especially under stress
- activates the immune system
- live yeast: 1000 x 10⁹ CFU/kg
- relieves the liver

Little helpers – great support

orgacell sc[®] is a premium complementary feedstuff consisting of a live and an inactive yeast component. Due to a low proportion of vegetable oil, the product is low in dust.

A well-functioning rumen provides the basis for a healthy, highyielding and fertile cow. Numerous trials demonstrate the positive effects on fiber digestion, performance and animal health. Therefore, yeasts deservedly have a firm place in animal nutrition.



Inactive yeast

Yeast cell walls have a special structure that gives them a high binding capacity. Thus, heavy metals, pathogens, and toxins can be bound and discharged from the animal without causing any harm. This not only supports the body's immune defenses, but also results in a higher level

of immunoglobulins in the colostrum, which are transferred directly to the newborn calf. In addition, inactive yeast has a supportive effect on the digestive tract, where it provides the microbes with nutrients and high-quality protein.

Live yeast

In **orgacell sc**[®], the live yeast *Saccharomyces cerevisiae* supports the rumen by binding O₂ there and stabilizing the pH value. This enables the beneficial bacteria to better digest fibers.

The high-quality microbial protein in the yeast makes more usable protein available to the cow, which has a positive effect on milk yield.

Feeding

orgacell sc[®] is mixed into the daily feed ration along with other dry components.

Dairy cattle: 40 g per animal per day
If the milk yield is high it makes sense to increase the dosage to 50 or 60 g per day.

Beef cattle: 70 g per animal per day

Calves for rearing: 20 g per animal per day

Packaging: 20 kg bag



A practical test from the Czech Republic showed the following results

All measured values have improved significantly: Less lactate and more fatty acids indicate a reduction in the risk of acidosis and an optimization of rumen processes. At the same time, the increase in fatty acids ensures a higher energy yield from the feed. Together with the increased glucose content, this leads to more milk in the tank. The beta-Hydroxybutyric acid value (BHBA in the blood) provides information about the energy balance. The higher it is, the more critical the ketosis risk. The low value shows that the live yeast has relieved the cow's metabolism.

	Untreated	Live yeast
pH value	6.4	6.6
lactate	16.4	9.3
fatty acids	104.7	112.0
Ø kg of milk ECM	37.0	39.4
glucose	3.2	3.3
BHBA	0.6	0.5

Sieve test

Using the simplest of means, the sieve test provides the easiest way to see the effects of feeding live yeast. Put a sample of manure in a common kitchen sieve and rinse until the water runs clear.

The undigested feed components will remain in the sieve. The amount and type of the residue shows the digestion's intensity.

After 3 - 4 weeks of feeding **orgacell sc**[®], repeat the test.

Through the feeding of live yeast the amount of residues is significantly reduced – especially the amount of undigested maize kernels is significantly reduced.



feed ration **without** live yeast




feed ration **with** live yeast

Effects of the live yeast *Saccharomyces cerevisiae* on the rumen

Consumes ruminal oxygen


Oxygen is toxic for the most significant ruminal microorganisms. Live yeast reduces oxygen, so the number of cellulose-degrading microorganisms increases. This may be observed in the animals' manure after just a short period of time (see pictures on the left): fibre and kernel residue is reduced. As live yeast binds the oxygen, a higher

amount of free hydrogen will be available for the formation of propionic acid. In the liver, this acid is subsequently transformed into the energy source glucose. In the dry period as well as during lactation, feeding low-energy rations results in an increased production of propionic acid in the rumen.



source: Lesaffre Feed Additives

Stabilizes the rumen pH value



Lactic acid-consuming bacteria are particularly stimulated and their ruminal population increases significantly.

An increased transformation from lactic acid into propionic acid further diminishes the risk of acidosis when feeding high-energy rations.

In high-energy rations, stabilizing the rumen pH is of special significance (see chart). If the milk yield is low, a high-fibre feed ration will satisfy the cow's energy requirements (see upper curve).

A rise in the level of performance requires high-energy rations containing starchy feedstuff / concentrate. These result in an increased production of lactic acid in the rumen and a subsequent lowering of the


pH when starch/carbohydrates are degraded (see lower curve). If the pH in the rumen drops below 5.8, the rumen villi are in danger of being irreversibly damaged by the acid and at the same time many bacteria die.

The middle curve shows how the live yeast stabilizes the rumen pH despite high milk yield.

Protecting the bacteria

A well-functioning rumen is the basic prerequisite for a healthy, high-performing cow. The microorganisms in the rumen work anaerobically. The more microorganisms are active in the rumen, the better the feed conversion. Consequently, the goal of every feeding must be to provide an optimal rumen environment for the bacteria.

orgacell sc® supports the rumen microbes and increases microbial activity. This allows the feed to be converted more effectively and also the feed intake increases.



Please note: detoxification function ceases!

If the pH drops below 6, a vital function of the rumen will falter: The degradation of toxins by single cell organisms like protozoa. Protozoa degrade complicated molecules such as mycotoxins but require a higher pH level for maintaining their vital functions. Thus, a rumen with frequently low pH levels bears the high risk that toxins are not degraded; and further on in the intestinal tract, will get into all organs via the bloodstream.


Double protection against toxins

orgacell sc® stabilizes the rumen pH and thus keeps the rumen villi healthy. These help to render possible toxins harmless. In addition, the yeast cell walls bind toxins and pathogenic germs, which are then discharged from the animal without causing harm.

A high-capacity rumen features a dense "lawn" of villi. Low pH values can really burn the rumen villi due to acid, so that the "lawn" gets holes and feed digestion deteriorates.

Sum of effects

Live yeast



Live yeast is activated by moisture and warmth. To guarantee this activation only in the rumen and not already on the feed table, our yeast is equipped with a protective coating.

In the rumen itself, live yeast has many effects:

- more effective fiber utilization
- less lactic acid stabilizes the rumen pH
- more propionic acid provides more energy
- relief from (heat) stress
- increase of rumen microbes due to O₂ binding
- increase of colostrum quality

Inactive yeast




Inactive yeast is also more than just a protein supplier:

- trace elements organically bound to amino acids
- B-vitamins and biotin
- binds toxins and pathogens
- stimulates the immune defenses

Each yeast source has its advantages, but both together form an unbeatable team:

- feed intake is increased
- animal health is improved
- animal can perform better



Distributor:

Tel.: +49 4262 - 20 74 -913 ▪ Fax: +49 4262 - 300 98 19 ▪ int@jbs.gmbh ▪ www.jbs.gmbh
joachim behrens scheessel gmbh ▪ Celler Straße 60 ▪ D-27374 Visselhövede